


syngenta.

SUBDUE MAXX is one of the best preventative and early curative systemic fungicides for effective control of Pythium in your greens.

What SUBDUE MAXX offers you:

- A highly effective, economical solution for Pythium in all turf species
- Protection of germinating seedlings during overseeding with high turf safety
- The combination of fast absorption and even systemic movement as well as contact action in soil, resulting in full plant protection
- MAXX technology allows for excellent tank mix compatibility with other fungicides and foliar fertilisers
- Low odour, non-staining formulation





To achieve great results, accurate application is important.

| Target | Syngenta Nozzle | Optimal Water Rate | Optimal Pressure Range |
|------------------|-----------------|--------------------|------------------------|
| Leaf and crown | XC 04 or XC 025 | 300 to 500 L/ha | 2 to 4 bar |
| Root and soil So | XC 08 | ~1000 L/ha | 2 to 4 bar |

SUBDUE MAXX is an excellent choice to use preventatively or early curatively during high infection risk periods. The fast absorption and systemic movement combines to ensure true "stopping power" for early curative Pythium Leaf Blight situations. For Pythium Root Rot a preventative plan is the only safe approach. SUBDUE MAXX is rapid and responsive treating problems before they are actually seen. Making a good start to the season should see turf through summer without a problem. Remember though that Pythium control depends heavily upon regular population control throughout the summer. Consecutive days of rain and high temperatures (25-40°C) will quickly advance Pythium populations.

Example: A sound strategy is therefore an early start with continued action.

- Late September SUBDUE MAXX @ 3 to 3.5 L/ha
- Late October SUBDUE MAXX @ 3 to 3.5 L/ha
- Late November propamocarb (600 g/L) @ 5 L/ha
- Before Christmas HEADWAY MAXX @ 9 L/ha
- Late January fosetyl (800 g/kg) @ 12.5 kg/ha


Give all applications a light wash in within 1 hour after spraying (6 to 10 mm).

| Target Diseases | Rate | SpraySure |
|--|-----------------|-----------|
| Pythium Leaf Bight | 1.7 to 3.5 L/ha | F |
| Pythium Root Rot, Seedling Damping Off | 1.7 to 3.5 L/ha | So |

Refer to the registered label for full details and instructions for use. F = Foliar applied So = Soil applied Su = Surfactant required







Active Ingredient

SUBDUE MAXX contains 240 g/L metalaxyl-M, a member of the phenylamide fungicides

GROUP

4 FUNGICIDE

Formulation and Packaging

1 L pack size

A micro emulsion concentrate (MEC) with minimal odour as the product does not contain petroleum solvents.

SUBDUE MAXX mixes completely with water and may be tank mixed with many commonly used fungicides and liquid fertilisers.

SUBDUE MAXX can be tank mixed with BANNER MAXX, HEADWAY MAXX, HERITAGE MAXX and PRIMO MAXX.



www.greencast.com.au



syngenta.